

AD-A139 237

REMARKS ON THE AMBROSETTI-PRODI PROBLEM(U) WISCONSIN  
UNIV-MADISON MATHEMATICS RESEARCH CENTER  
D G COSTA ET AL. JAN 84 MRC-TSR-2633 DAAG29-80-C-0041

1/1

UNCLASSIFIED

F/G 12/1

NL


END

DATE

FILED

4-24

DTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD A139237

MRC Technical Summary Report # 2633

REMARKS ON THE AMBROSETTI-PRODI PROBLEM

D. G. Costa and D. G. de Figueiredo

**Mathematics Research Center  
University of Wisconsin—Madison  
610 Walnut Street  
Madison, Wisconsin 53705**

January 1984

(Received November 21, 1983)

DTIC  
ELECTE  
MAR 22 1984  
S B D

**Approved for public release  
Distribution unlimited**

Sponsored by

U. S. Army Research Office  
P. O. Box 12211  
Research Triangle Park  
North Carolina 27709

**DTIC FILE COPY**

4

037

UNIVERSITY OF WISCONSIN-MADISON  
MATHEMATICS RESEARCH CENTER

REMARKS ON THE AMBROSETTI-PRODI PROBLEM

D. G. Costa\* and D. G. de Figueiredo\*\*

Technical Summary Report # 2633

January 1984

ABSTRACT

✓ The Ambrosetti-Prodi boundary value problem with an asymptotically linear nonlinearity is considered. Under general conditions on the nonlinearity it is shown that there exist positive and negative solutions. In the case when the domain is a ball in  $\mathbb{R}^n$  and the nonlinearity crosses the first  $n$  eigenvalues, corresponding to radial eigenfunctions, it is proved that there are at least  $n + 1$  radial solutions.

AMS (MOS) Subject Classification: 35J65, 47H15.

Key Words: Ambrosetti-Prodi problem, positive and negative solutions, radial solutions

Work Unit Number 1 (Applied Analysis)

\*Universidade de Brasilia (Brazil). Partially supported by CNPq/Brazil.

\*\*Universidade de Brasilia (Brazil) and Guggenheim Fellow (1983).

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041.

### SIGNIFICANCE AND EXPLANATION

Let  $\Omega \subset \mathbb{R}^N$  be a bounded smooth domain. Let  $\lambda_1$  be the first eigenvalue of  $-\Delta$  on  $H_0^1(\Omega)$  and let  $\phi_1 > 0$  be a corresponding eigenfunction. The Ambrosetti-Prodi problem

$$(1)_t \quad -\Delta u = f(x, u) + h(x) + t\phi_1(x) \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega,$$

where  $t \in \mathbb{R}$ ,  $h \in C(\bar{\Omega})$ ,  $\int h \phi_1 = 0$ , is studied. First it is proved that if

$\lim_{s \rightarrow -\infty} f(x,s)/s < \lambda_1 < \lim_{s \rightarrow +\infty} f(x,s)/s$  then problem  $(1)_t$  has a large positive and a large negative solution for  $t < 0$  large. The interest in this type of solution comes from the fact that they play a special role in the existence of other solutions. This has been observed before for special classes of problem  $(1)_t$  by Lazer-McKenna, Solimini and Ambrosetti. If the limit of  $f(x,s)/s$  at  $+\infty$  is larger than a higher eigenvalue, it is expected that other solutions, besides the positive and negative ones, will appear. Previous results of Hofer, Solimini and Ambrosetti indicate that this is the case. They have been able to prove existence of up to six solutions. A connection between the number of solutions and the number of eigenvalues which are crossed has been proved to exist in the o.d.e. case by Lazer-McKenna. Here the authors consider the case of a ball in  $\mathbb{R}^N$  and prove the existence of many radial solutions. These solutions are characterized by the number of their "nodal lines".



**The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the authors of this report.**

les  
r

Dist Social

A-1

# REMARKS ON THE AMBROSETTI-PRODI PROBLEM

D. G. Costa\* and D. G. de Figueiredo\*\*

Let  $\Omega \subset \mathbb{R}^N$  be a bounded smooth domain. Let  $\lambda_1 < \lambda_2 < \dots < \lambda_k < \dots$  denote the eigenvalues of  $-\Delta$  on  $\Omega$  with Dirichlet boundary condition and  $\phi_1 > 0$  an eigenfunction corresponding to  $\lambda_1$ . We consider the Ambrosetti-Prodi problem

$$(1)_t \quad -\Delta u = f(x, u) + h(x) + t\phi_1(x) \text{ in } \Omega, u = 0 \text{ on } \partial\Omega,$$

where  $h \in C(\bar{\Omega})$ ,  $\int \phi_1 = 0$ , and  $f: \bar{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$  is a continuous function such that there exist the limits

$$(2) \quad \lim_{s \rightarrow \pm\infty} \frac{f(x, s)}{s} = f_{\pm} \text{ uniformly for } x \in \bar{\Omega}, \text{ with}$$

$$(3) \quad f_- < \lambda_1 < f_+, \quad f_+ \neq \lambda_j.$$

We assume without loss of generality that  $f(x, 0) \equiv 0$ .

In Section 1 of the present paper we shall show that for  $t < 0$  sufficiently large problem  $(1)_t$  possesses a large positive solution and a large negative solution. We remark that for a special class of problems  $(1)_t$  existence of a negative solution was first observed by Lazer-McKenna [3]. Subsequently, Ambrosetti [1] and Solimini [5], again for a special class of problems  $(1)_t$  proved the existence of both a negative and a positive solution for  $t < 0$  sufficiently large. In [2], one of the authors proved existence of a negative solution for a general class of Ambrosetti-Prodi problems, including the superlinear case. At present we do not know whether a positive solution exists in the case of nonlinearities  $f$  which grow more rapidly than linear.

In Section 2 we consider the case when  $\Omega$  is a ball and  $f$  is a  $C^1$  function which does not depend explicitly on  $n$ . Let us denote by  $\mu_1 < \mu_2 < \dots$  the eigenvalues

---

\*Universidade de Brasília (Brazil). Partially supported by CNPq/Brazil.

\*\*Universidade de Brasília (Brazil) and Guggenheim Fellow (1983).

---

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041.

of  $-\Delta$  acting on the radial functions of  $H_0^1(\Omega)$  (It is well-known that  $\mu_1 = \lambda_1$ ). We assume there that the limits below exist and satisfy

$$(4) \quad f_- \equiv \lim_{s \rightarrow -\infty} f'(s) < \mu_1, \quad \mu_n < f_+ \equiv \lim_{s \rightarrow +\infty} f'(s) < \mu_{n+1}, \quad n > 1.$$

It is then proved that for  $t < 0$  sufficiently large  $(1)_t$  possesses a radial positive solution  $U_t$  and a radial negative solution  $u_t$ . It is also proved that given any  $c \neq 0$  there exists a radial solution  $u$  of the equation in  $(1)_t$  such that  $u - c \in H_0^1(\Omega)$  and  $u - U_t$  has at least  $n$  "nodal lines". More precisely there are at least  $n$  concentric spheres where the function  $u - U_t$  vanishes.

In Section 3 we prove under the same assumptions of the previous section that problem  $(1)_t$  has at least  $n + 1$  radial solutions for  $t < 0$  sufficiently large. Here we use a shooting argument in a manner similar to the work of Lazer-McKenna [4]. In their paper they treated a two-point boundary value problem, which is the o.d.e. analogue of  $(1)_t$ , with  $-\Delta u$  replaced by  $-u$ . Our analysis would correspond in their case to looking for solutions which are symmetric with respect to the middle point of the interval. Without this assumption of symmetry they can actually obtain  $2n$  solutions. We believe that this is also true in the p.d.e. case because we are in fact crossing also a number of eigenvalues that correspond to non-radial eigenfunctions. As a matter of fact we prove in Section 4 that this is the case when  $N = 3$ . Finally we observe that in the case when  $\Omega$  is a region between two concentric spheres there are at least  $2n$  solutions for problem  $(1)_t$ ,  $t < 0$  large.

#### 1. Existence of large positive and negative solutions

In this section we let  $\Omega \subset \mathbb{R}^N$  be an arbitrary bounded smooth domain. Let  $\lambda_1 < \lambda_2 < \dots < \lambda_k < \dots$  denote the eigenvalues of  $-\Delta$  on  $\Omega$  with Dirichlet boundary condition and  $\phi_1 > 0$  an eigenfunction corresponding to  $\lambda_1$ . We consider the Ambrosetti-Prodi problem

$$(1)_t \quad -\Delta u = f(x, u) + h(x) + t\phi_1(x) \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega,$$

where  $h \in C(\bar{\Omega})$ ,  $\int h \phi_1 = 0$ , and  $f: \bar{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$  is a continuous function such that

$f(x, 0) \equiv 0$  and there exist the limits

$$(2) \quad \lim_{s \rightarrow \pm\infty} \frac{f(x, s)}{s} = f_{\pm}, \text{ uniformly for } x \in \bar{\Omega}, \text{ with}$$

$$(3) \quad f_- < \lambda_1 < f_+, \quad f_+ \neq \lambda_j.$$

We shall show that for  $t < 0$  sufficiently large problem  $(1)_t$  possesses a large positive solution and a large negative solution. To that end consider the related problems

$$(1)_{\alpha}^{\pm} \quad -\Delta v = f_{\pm} v + F(x, v + \alpha \phi_1) + h \text{ in } \Omega, \quad v = 0 \text{ on } \partial\Omega, \quad (\alpha \in \mathbb{R}),$$

where  $F(x, s) = f(x, s) - (f_+ s^+ - f_- s^-)$ , so that

$$(4) \quad \lim_{|s| \rightarrow \infty} \frac{F(x, s)}{s} = 0 \text{ uniformly for } x \in \bar{\Omega}.$$

Denote by  $S^{\pm}(\alpha)$  the set of solutions of  $(1)_{\alpha}^{\pm}$  and let  $M^{\pm}(\alpha) = \sup \{ \|v\|_{C^1} : v \in S^{\pm}(\alpha) \}$ .

Lemma 1.  $\lim_{|\alpha| \rightarrow \infty} \frac{M^{\pm}(\alpha)}{\alpha} = 0.$

Proof. We shall consider only the case of  $M^+(\alpha)$ , the other case being entirely similar. Any solution  $v \in S^+(\alpha)$  is a solution of the equation  $v = K_+ [F(\cdot, v + \alpha \phi_1) + h]$ , where  $K_+: L^2(\Omega) \rightarrow L^2(\Omega)$  is the resolvent operator  $K_+ = (-\Delta - f_+)^{-1}$  (which exists in view of (3)). As it is well-known,  $K_+: C^0(\bar{\Omega}) \rightarrow C^1(\bar{\Omega})$  is a bounded operator and, therefore, any  $v \in S^+(\alpha)$  satisfies

$$\|v\|_{C^1} \leq c_+ \|F(\cdot, v + \alpha \phi_1) + h\|_{C^0},$$

or yet, since one has by (4) that  $|F(x, s)| \leq \varepsilon |s| + c_{\varepsilon}$  for all  $x \in \bar{\Omega}$ ,  $s \in \mathbb{R}$ , where  $\varepsilon > 0$  can be chosen arbitrarily (and  $c_{\varepsilon}$  depends only on  $\varepsilon$ ),

$$\|v\|_{C^1} \leq c_+ [\varepsilon \|v\|_{C^0} + \varepsilon |\alpha| \|\phi_1\|_{C^0} + c_{\varepsilon} + \|h\|_{C^0}].$$

Choosing  $\varepsilon > 0$  so that  $1 - c_+ \varepsilon > 1/2$  one obtains

$$\|v\|_{C^1} \leq 2 c_+ \varepsilon |\alpha| \|\phi_1\|_{C^0} + 2 c_+ (c_{\varepsilon} + \|h\|_{C^0}),$$

from which the claim follows readily. ■

Theorem 1. Suppose that (2) and (3) hold. Then there exists  $t_0 < 0$  such that for all  $t < t_0$  problem  $(1)_t$  has a positive solution  $u_t$  and a negative solution  $u_t$  satisfying

$$(5) \quad \lim_{t \rightarrow -\infty} \left\| \frac{u_t}{t} - \frac{\phi_1}{\lambda_1 - f_+} \right\|_{C^1} = \lim_{t \rightarrow -\infty} \left\| \frac{u_t}{t} - \frac{\phi_1}{\lambda_1 - f_-} \right\|_{C^1} = 0.$$

Proof. Let  $\eta > 0$  be such that  $w + \phi_1 > 0$  for all  $w$  satisfying

$\|w\|_{C^1} < \eta$ ,  $w|_{\partial\Omega} = 0$ . By Lemma 1 there exists  $\alpha_0 > 0$  such that  $M^+(\alpha) < \eta\alpha$  for all  $\alpha > \alpha_0$ , that is,

$$(6) \quad \|v\|_{C^1} < \eta\alpha$$

for any solution  $v$  of  $(1)_{\alpha}^+$ ,  $\alpha > \alpha_0$ . For any such  $v$ , in view of the choice of  $\eta$  and the fact that  $\|v/\alpha\|_{C^1} < \eta$ , we have that

$$(7) \quad U \equiv v + \alpha\phi_1 = \alpha\left(\frac{v}{\alpha} + \phi_1\right) > 0.$$

So  $U$  is a positive solution of the problem

$$-\Delta u = f_+(u - \alpha\phi_1) + \xi(x, u) + h + \alpha\lambda_1\phi_1 \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega,$$

for  $\alpha > \alpha_0$ , that is,  $U$  is a positive solution of  $(1)_t$  for

$t = \alpha(\lambda_1 - f_+) < \alpha_0(\lambda_1 - f_+) \equiv t_0^+$ . And, in view of (6) and (7),  $U/t$  satisfies the inequality

$$\left\| \frac{U}{t} - \frac{\phi_1}{\lambda_1 - f_+} \right\|_{C^1} = \left\| \frac{v}{t} \right\|_{C^1} < \frac{\eta}{f_+ - \lambda_1}.$$

Similarly, we prove the existence of a negative solution  $u_t$  of  $(1)_t$  for

$t < -\alpha_0(\lambda_1 - f_-) \equiv t_0^-$  satisfying

$$\left\| \frac{u_t}{t} - \frac{\phi_1}{\lambda_1 - f_-} \right\|_{C^1} < \frac{\eta}{\lambda_1 - f_-}.$$

The proof is complete by letting  $t_0 = \min\{t_0^+, t_0^-\}$  and noticing that the chosen  $\eta > 0$  can be taken arbitrarily small. ■

## 2. The case when $\Omega$ is a ball

We now let  $\Omega$  be the unit ball  $B = B_1(0) \subset \mathbb{R}^N$  and consider the Ambrosetti-Prodi problem

$$(8)_t \quad -\Delta u = f(u) + h + t\phi_1 \text{ in } B, \quad u = 0 \text{ on } \partial B,$$

where  $h \in C(\bar{B})$  is a given radial function with  $\int h \phi_1 = 0$  and  $f: \mathbb{R} \rightarrow \mathbb{R}$  is a  $C^1$  function satisfying  $f(0) = 0$ ,

$$(9) \quad \lim_{s \rightarrow -\infty} f'(s) \equiv f_- < \mu_1, \quad \mu_n < \lim_{s \rightarrow +\infty} f'(s) \equiv f_+ < \mu_{n+1}, \quad n \geq 1.$$

Here we are denoting by  $\mu_1 = \lambda_1 < \mu_2 < \dots$  the eigenvalues of  $-\Delta$  acting on the radial functions of  $H_0^1(B)$ , that is, the eigenvalues of the problem

$$(10) \quad -u'' - \frac{N-1}{r} u' \equiv \mu u, \quad 0 < r < 1, \quad u'(0) = u(1) = 0.$$

As it is well-known, these eigenvalues are given by  $\mu_j = v_j^2$  where the  $v_j$ 's are the positive zeros of the Bessel function  $J_{(N-2)/2}$ , and corresponding eigenfunctions are given by  $\phi_j(r) = r^{-(N-2)/2} J_{(N-2)/2}(v_j r)$ .

Theorem 2. Under assumption (9), there exists  $t_1 < 0$  such that for all  $t < t_1$  problem  $(8)_t$  has a radial positive solution  $U_t$  and a radial negative solution  $u_t$  satisfying (5).

Proof. In view of Theorem 1 there exists  $t_0 < 0$  such that for all  $t < t_0$  problem  $(8)_t$  has a positive solution  $U_t$  and a negative solution  $u_t$  satisfying (5). It remains to show that both  $U_t$  and  $u_t$  are radial functions for  $t < t_0$  sufficiently large. Indeed, letting  $z_t = \partial U_t / \partial \omega$  denote any angular derivative of  $U_t$ ,  $z_t$  satisfies  $-\Delta z_t = f'(U_t) z_t$  in  $B$ ,  $z_t = 0$  on  $\partial B$ . Therefore if we assume that  $z_t \not\equiv 0$ ,  $z_t$  is an eigenfunction of the above problem corresponding to the eigenvalue  $1 = \lambda_j(f'(U_t))$ . But then the fact that  $U_t(x) \rightarrow +\infty$  for all  $x \in B$  and the Lebesgue dominated convergence theorem imply that  $f'(U_t) \rightarrow f_+$  in  $L^p$ ,  $p > 1$ , and hence that  $1 = \lambda_j(f'(U_t)) \rightarrow \lambda_j(f_+) = \lambda_j/f_+$  as  $t \rightarrow -\infty$ , which is a contradiction. Similarly we show that  $u_t$  is a radial function for  $t < t_0$  sufficiently large. ■

Now we make the change of (dependent) variables  $u = w + U_t$  to rewrite problem  $(8)_t$  as

$$-\Delta w = f(w+U_t) - f(U_t) \text{ in } B, \quad w = 0 \text{ on } \partial B,$$

or yet as

$$(11)_t \quad -\Delta w = f_+ w + g(w+U_t) - g(U_t) \text{ in } B, \quad w = 0 \text{ on } \partial B,$$

where  $g(s) \equiv f(s) - f_+ s$ . Notice that, in view of (9),  $g: \mathbb{R} \rightarrow \mathbb{R}$  is a  $C^1$  function satisfying

$$(12) \quad \lim_{s \rightarrow -\infty} g'(s) = f_- - f_+, \quad \lim_{s \rightarrow +\infty} g'(s) = 0.$$

On the other hand, given  $0 \neq \bar{c} \in \mathbb{R}$ , let us define the function

$$\bar{v}(r) = \frac{\bar{c}}{r^{(N-2)/2}} \frac{J_{(N-2)/2}(\sqrt{f_+} r)}{J_{(N-2)/2}(\sqrt{f_+})}, \quad r = |x|.$$

Then it is easy to see that  $\bar{v}$  is a solution of the problem

$$(13) \quad -\Delta \bar{v} = f_+ \bar{v} \text{ in } B, \quad \bar{v} = \bar{c} \text{ on } \partial B,$$

and, since  $v_n < \sqrt{f_+} < v_{n+1}$  by (9),  $\bar{v}$  vanishes on the  $n$  concentric spheres  $r = r_j = v_j / \sqrt{f_+}$  ( $j=1, \dots, n$ ) in  $B$ .

Theorem 3. Suppose (9) holds and let  $0 \neq \bar{c} \in \mathbb{R}$  be given. Then there exists

$\bar{t} = \bar{t}(\bar{c}) < 0$  such that for all  $t < \bar{t}$  problem

$$(14)_{t, \bar{c}} \quad -\Delta u = f(u) + h + t\phi_1 \text{ in } B, \quad u = \bar{c} \text{ on } \partial B,$$

has a radial solution  $\bar{u}$  with the property that  $\bar{u} - U_t$  vanishes on  $n$  concentric spheres in  $B$ .

Proof. We want to find a radial solution  $w$  of  $(11)_t$ , with the boundary condition replaced by  $w = \bar{c}$  on  $\partial B$ , and such that  $w$  vanishes on  $n$  concentric spheres in  $B$ . Since the function  $\bar{v}$  has this latter property and its  $n$  zeros in  $(0, 1)$  (as a function of  $r$ ) are all simple, it suffices to find a solution  $z$  of the problem

$$(15)_t \quad -\Delta z = f_+ z + g(z+\bar{v}+U_t) - g(U_t) \text{ in } B, \quad z = 0 \text{ on } \partial B,$$

with  $\|z\|_{C^0} < \bar{\varepsilon}$ ,  $\bar{\varepsilon} = \bar{\varepsilon}(\bar{c})$  sufficiently small, for then  $\bar{u} = z + \bar{v} + U_t$  is a solution of

(14) with the desired property that  $\bar{U} - U_t = z + \bar{V}$  vanishes on  $n$  concentric spheres in  $B$ . (We point out that the same argument used in the proof of Theorem 2 shows that the solutions of  $(15)_t$  are radial for  $t < 0$  sufficiently large). So Lemma 2 below concludes the proof.

Lemma 2. Given  $\bar{\epsilon} > 0$  there exists  $\bar{t} = \bar{t}(\bar{\epsilon})$  such that for all  $t < \bar{t}$  problem  $(15)_t$  has a unique solution  $z$  satisfying  $\|z\|_{C^0} < \bar{\epsilon}$ .

Proof. We observe that solving  $(15)_t$  is equivalent to solving the operator equation

$$z = KG_t(z) \equiv T_t(z),$$

where  $K : L^2(B) \rightarrow H^1_0(B)$  is the resolvent operator  $K = (-\Delta - f_+)^{-1}$  and  $G_t$  is the Nemytskii operator associated with the function  $g(z + \bar{V}(r) + U_t(r)) - g(U_t(r))$ . We also observe that the regularity theory for elliptic equations implies that  $K$  maps  $L^p(B)$  continuously into  $W^{2,p}(B)$  and (12) implies that  $G_t$  maps  $L^p(B)$  continuously into itself, so that  $T_t$  maps  $L^p(B)$  into  $W^{2,p}(B)$ . Given the ball  $B_{\frac{\epsilon}{2}}(0)$  in  $C^0(\bar{B})$ , we shall show that for  $t < 0$  sufficiently large we have  $T_t(B_{\frac{\epsilon}{2}}(0)) \subset B_{\frac{\epsilon}{2}}(0)$  and  $T_t$  is a contraction. Indeed, fixing  $p > N/2$  and using the Sobolev imbedding theorem, we have

$$\|T_t(t)\|_{C^0} < c\|T_t(z)\|_{W^{2,p}} = c\|KG_t(t)\|_{W^{2,p}} < \text{const.} \|G_t(z)\|_{L^p},$$

and  $\|G_t(z)\|_{L^p} \rightarrow 0$  as  $t \rightarrow -\infty$  uniformly for  $z \in B_{\frac{\epsilon}{2}}(0)$ , since we can estimate

$$\|G_t(z)\|_{L^p} < \delta(t)\|z + \bar{V}\|_{C^0} \quad \text{where } \delta(t) \rightarrow 0 \text{ as } t \rightarrow -\infty \quad (\text{In computing } \|G_t(z)\|_{L^p} \text{ apply}$$

the mean value theorem for the function  $g$  and use the Lebesgue dominated convergence theorem, keeping in mind that  $z + \bar{V} + U_t \rightarrow +\infty$  pointwisely in  $B$  and that

$g'(s) \rightarrow 0$  as  $s \rightarrow +\infty$ ). Therefore,  $T_t(B_{\frac{\epsilon}{2}}(0)) \subset B_{\frac{\epsilon}{2}}(0)$  for  $t < 0$  sufficiently large.

Similarly, for arbitrary  $z_1, z_2 \in B_{\frac{\epsilon}{2}}(0)$ , we have the estimate

$$\|T_t(z_1) - T_t(z_2)\|_{C^0} < \text{const.} \|G_t(z_1) - G_t(z_2)\|_{L^p} < \text{const.} \delta(t)\|z_1 - z_2\|_{C^0},$$

where  $\delta(t) \rightarrow 0$  as  $t \rightarrow -\infty$ , so that  $T_t : B_{\frac{\epsilon}{2}}(0) \rightarrow B_{\frac{\epsilon}{2}}(0)$  is a contraction for  $t < 0$  sufficiently large. The proof is complete. ■

Remark. It should be pointed out that we stated and proved Theorem 3 as a p.d.e. result, ignoring for the time being its natural 1-dimensional character (i.e., search of radial solutions in a ball). The reason for that is the fact that indeed, for an arbitrary bounded smooth domain  $\Omega$ , Theorem 3 has the following analogue which we now describe.

For given  $0 \neq c \in \mathbb{R}$ , let  $v_c$  denote the solution of the problem  $-\Delta v = f_+ v$  in  $\Omega$ ,  $v = c$  on  $\partial\Omega$  (i.e.,  $v_c = (-\Delta)(-\Delta - f_+)^{-1}c$ ). Suppose that for some  $\bar{c} \neq 0$ ,  $\bar{v} = \frac{v}{\bar{c}}$  satisfies the condition that  $\nabla \bar{v}(x) \neq 0$  whenever  $\bar{v}(x) = 0$ , that is, the graph of  $\bar{v}$  is transversal to  $\Omega \times \{0\}$  at the points of  $M \times \{0\}$ , where  $M = \{x \in \Omega \mid \bar{v}(x) = 0\}$  is the "null manifold" of  $\bar{v}$ . Let  $m$  be the number of components of  $M$ . Then, for  $t < 0$  sufficiently large (and under hypotheses (9)) there exists a solution  $u$  of the problem

$$-\Delta u = f(u) + h + t\phi_1 \text{ in } \Omega, \quad u = c \text{ on } \partial\Omega,$$

such that  $u - U_t$  has at least  $m$  "nodal lines", that is, the null manifold of  $u - U_t$  has at least  $m$  components.

### 3. Existence of many solutions

In this section we consider again the case when  $\Omega$  is the unit ball  $B = B_1(0) \subset \mathbb{R}^N$  and prove under the same assumption (9) of the previous section that the Ambrosetti-Prodi problem  $(8)_t$  has at least  $n + 1$  radial solutions for  $t < 0$  sufficiently large. For that matter we shall use a shooting argument as in [5].

It should be remarked that only from now on is that the one-dimensional character of problem  $(8)_t$  will play an important role in proving the existence of other solutions besides  $U_t > 0$  and  $u_t < 0$ , when  $n \geq 2$  in (9).

So, we start by rewriting  $(8)_t$  as

$$(16)_t \quad -u'' - \frac{N-1}{r} u' = f(u) + h + t\phi_1, \quad 0 < r < 1, \quad u'(0) = u(1) = 0,$$

which in turn, again through the change of dependent variables  $u(r) = w(r) + U_t(r)$ , can be rewritten as

$$(17a)_t \quad -w'' - \frac{N-1}{r} w' = f_+ w + g(w + U_t) - g(U_t), \quad 0 < r < 1,$$

$$(17b) \quad w'(0) = w(1) = 0,$$

where we recall that  $g(s) \equiv f(s) - f_+ s$ .

Lemma 3. The initial value problem

$$-w'' - \frac{N-1}{r} w' = f_+ w + g(w + U_t(r)) - g(U_t(r)), \quad 0 < r < 1, \quad w(0) = a, \quad w'(0) = 0,$$

has a unique solution  $w(r) = w(r; a)$  which is defined for  $0 < r < 1$  and depends continuously on  $a \in \mathbb{R}$ .

Proof. Since  $-w'' - \frac{N-1}{r} w' = -\frac{1}{r^{N-1}} (r^{N-1} w')'$ , it can be seen that the initial value problem in question is equivalent to the following Volterra integral equation

$$w(r) = a + \int_0^r K(r, \sigma) F(\sigma, w(\sigma)) d\sigma,$$

where  $F(\sigma, s) \equiv f_+ s + g(s + U_t(\sigma)) - g(U_t(\sigma))$  and  $K(r, \sigma) = \frac{\sigma}{N-2} \left[ \left( \frac{\sigma}{r} \right)^{N-2} - 1 \right]$ . Since the kernel  $K(r, \sigma)$  is nice, the result follows from the standard theory of Volterra equations. ■

Theorem 4. Under assumption (9), there exists  $t_2 < 0$  such that for all  $t < t_2$  problem (17)<sub>t</sub> has at least  $n$  distinct non-trivial solutions  $w_0, w_1, \dots, w_{n-1}$  with the property that  $w_j(r)$  has exactly  $j$  simple zeros in the open interval  $(0, 1)$  and  $w_j(0) < 0$ ,  $j = 0, \dots, n-1$ . (Therefore, for  $t < t_2$  problem (8)<sub>t</sub> has at least the  $n+1$  distinct radial solutions  $U_t, w_j + U_t$ ,  $j = 0, \dots, n-1$ .)

Proof. From Theorem 2 we already know 2 solutions for problem (17)<sub>t</sub>, namely  $w \equiv 0$  (corresponding to the positive solution  $U_t$  of (8)<sub>t</sub>) and  $w_0 = u_t - U_t < 0$  (corresponding to the negative solution  $u_t$  of (8)<sub>t</sub>), provided  $t < t_1$ . On the other hand, from Theorem 3 we have a solution  $\bar{w} = \bar{u} - U_t$  of equation (17a)<sub>t</sub>, for  $t < \bar{t}$ , such that  $\bar{w}(r)$  has  $n$  simple zeros in the open interval  $(0, 1)$  and  $\bar{w}(0) < 0$  (take  $\bar{c}$  in (14)<sub>t, \bar{c}}</sub> so that the

function  $\bar{v}$  in (13) satisfies  $\bar{v}(0) < 0$ . Therefore, in view of Lemma 3, we can use the idea of Lemma 2.3 in [4] to show that, for each  $0 < j < n$ , problem  $(17)_t$  has a solution  $w_j(r)$  with exactly  $j$  simple zeros in  $(0,1)$  and such that  $w_j(0) < 0$ . ■

#### 4. The case $N = 3$

In the case that  $\Omega$  is the unit ball  $B = B_1(0)$  in  $\mathbb{R}^3$  we are able to improve Theorem 4 and obtain  $2n$  radial solutions for the Ambrosetti-Prodi problem  $(8)_t$ ,  $t < 0$  sufficiently large. Namely, we have

Theorem 5. Let  $N = 3$  and suppose (9) holds. Then, for all  $t < 0$  sufficiently large, problem  $(17)_t$  has at least  $2n-1$  distinct non-trivial solutions  $w_j$ ,  $j = 0, \dots, n-1$ , and  $\tilde{w}_j$ ,  $j = 1, \dots, n-1$ , with the property that  $w_j$  and  $\tilde{w}_j$  have exactly  $j$  simple zeros in  $(0,1)$  and  $w_j(0) < 0$ ,  $\tilde{w}_j(0) > 0$ . (Therefore, for  $t < 0$  sufficiently large, problem  $(8)_t$  has at least the following  $2n$  distinct radial solutions:  $U_t$ ,  $w_j + U_t$ ,  $j = 0, \dots, n-1$ ,  $\tilde{w}_j + U_t$ ,  $j = 1, \dots, n-1$ .)

Proof. By making the change of variables  $v(r) = rw(r)$  and letting  $V_t(r) \equiv rU_t(r)$ , we transform  $(17)_t$  into the problem

$$(18)_t \quad -v'' = f_+ v + \tilde{g}(r, v + V_t) - \tilde{g}(V_t), \quad 0 < r < 1, \quad v(0) = v(1) = 0,$$

where  $\tilde{g}(r, v) \equiv rg(v/r)$ . Notice that (9) (hence (12)) implies that  $\tilde{g}$  is continuous on  $[0,1] \times \mathbb{R}$  (by defining  $\tilde{g}(0, v) = (f_+ - f_-)v$ ),  $\tilde{g}$  is of class  $C^1$  on  $(0,1) \times \mathbb{R}$ ,  $\tilde{g}_v$  is continuous on  $[0,1] \times (\mathbb{R} \setminus \{0\})$  and satisfies  $\lim_{v \rightarrow \pm\infty} \tilde{g}_v(r, v) = f_- - f_+$ .

$\lim_{v \rightarrow \pm\infty} \tilde{g}_v(r, v) = 0$  uniformly for  $r \in [0,1]$ . Therefore, problem  $(18)_t$  can be treated in a similar manner as the problem in [4]. It then follows that  $(18)_t$  has solutions  $v_j$ ,  $j = 0, \dots, n-1$ , and  $\tilde{v}_j$ ,  $j = 1, \dots, n-1$ , such that  $v_j$  and  $\tilde{v}_j$  have exactly  $j$  simple zeros in  $(0,1)$  and  $v_j'(0) < 0$ ,  $\tilde{v}_j'(0) > 0$ . Going back to our original (dependent) variable we first claim that if  $v(r)$  is a solution of  $(18)_t$  then  $w(r) = v(r)/r$  is a solution of  $(17)_t$ . Indeed, the only non-obvious property to check is that

$w(r)$  satisfies the boundary condition  $w'(0) = 0$ . For that we observe that any solution  $v(r)$  of  $(18)_t$  satisfies  $v''(0) = 0$ , so that using l'Hospital's rule we obtain

$$\lim_{r \rightarrow 0} w'(r) = \lim_{r \rightarrow 0} \frac{rw'(r) - v(r)}{r^2} = \lim_{r \rightarrow 0} \frac{r v''(r)}{2r} = \frac{v''(0)}{2} = 0.$$

Consequently  $w_j(r) = v_j(r)/r$  and  $\tilde{w}_j(r) = \tilde{v}_j(r)/r$  are solutions of  $(17)_t$  with the stated properties. ■

Remark. We observe that in the case  $\Omega$  is the region between two concentric spheres in  $\mathbb{R}^N$ , say  $\varepsilon < r < 1$ , we again obtain  $2n$  radial solutions for the corresponding Ambrosetti-Prodi problem. Indeed, we are led to the o.d.e. problem  $(17a)_t$ ,

$w(\varepsilon) = w(1) = 0$ , or yet, making the change of variable  $v(r) = r^{(N-1)/2} w(r)$ , to the self-adjoint problem

$$-v'' + \frac{C_N}{r^2} v = f_+ v + \tilde{g}(r, v + V_t) - \tilde{g}(r, V_t), \quad \varepsilon < r < 1, \quad v(\varepsilon) = v(1) = 0,$$

where  $C_N = (N-1)(N-3)/4$ , and  $V_t(r)$  and  $\tilde{g}(r, v)$  are as before.

#### REFERENCES

- [1] Ambrosetti, A. - "Elliptic equations with jumping nonlinearities", to appear.
- [2] de Figueiredo, D. G. - "On the superlinear Ambrosetti-Prodi problem", MRC Tech. Rep. #2522 (May 1983).
- [3] Lazer, A. C. and McKenna, P. J. - "On the number of solutions of a nonlinear Dirichlet problem", J. Math. Anal. Appl. 84 (1981), 282-294.
- [4] Lazer, A. C. and McKenna, P. J. - "On a conjecture related to the number of solutions of a nonlinear Dirichlet problem", to appear.
- [5] Solimini, S. - "Existence of a third solution for a class of b.v.p. with jumping nonlinearities", Nonlinear Anal. TMA 7 n. 8 (1983), 917-927.

DGC/DGF;jgb

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2633	2. GOVT ACCESSION NO. AD-A139237	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) REMARKS ON THE AMBROSETTI-PRODI PROBLEM		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) D. G. Costa and D. G. de Figueiredo		8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 1 - Applied Analysis
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P.O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE January 1984
		13. NUMBER OF PAGES 12
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Ambrosetti-Prodi problem, positive and negative solutions, Radial solutions		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  The Ambrosetti-Prodi boundary value problem with an asymptotically linear nonlinearity is considered. Under general conditions on the nonlinearity it is shown that there exist positive and negative solutions. In the case when the domain is a ball in $\mathbb{R}^n$ and the nonlinearity "crosses" the first $n$ eigenvalues, corresponding to radial eigenfunctions, it is proved that there are at least $n + 1$ radial solutions.		